



UNIVERSITI PUTRA MALAYSIA

**EFFECTS OF WASHING WITH POTASSIUM AND SODIUM
HYDROXIDE MIXTURES IN THE DEVELOPMENT OF
RANCIDITY IN BLACK TILAPIA (*OREOCHROMIS
MOSSAMBICUS*) FILLETS AND THE FRYING OIL**

ANIDA YUSOFF

FSMB 2003 37

**EFFECTS OF WASHING WITH POTASSIUM AND SODIUM HYDROXIDE
MIXTURES IN THE DEVELOPMENT OF RANCIDITY IN BLACK TILAPIA
(*OREOCHROMIS MOSSAMBICUS*) FILLETS AND THE FRYING OIL**

By

ANIDA YUSOFF

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
In Fulfilment of the Requirement for the Degree of Master of Science**

January 2003



Dedicated to my beloved.....

Parents ...mak, abah..

SistersAlong, Nyah...

Twin sisterYa

and

Brother...Aliff

Abstract of the thesis presented to the Senate of the Universiti Putra Malaysia in
fulfilment of the requirements for the degree of Master of Science

**EFFECTS OF WASHING WITH POTASSIUM AND SODIUM HYDROXIDE
MIXTURES IN THE DEVELOPMENT OF RANCIDITY IN BLACK TILAPIA
(*OREOCHROMIS MOSSAMBICUS*) FILLETS AND THE FRYING OIL**

By

ANIDA YUSOFF

January 2003

Chairman : Associate Professor Dr. Jamilah Bakar, Ph.D.

Faculty : Food Science and Biotechnology

A study was carried out to determine the effectiveness of potassium hydroxide (KOH) and sodium hydroxide (NaOH) in removing the earthy flavor and odor in black tilapia (*Oreochromis mossambicus*) muscle. Four different mixtures of KOH and NaOH were also prepared, each in three different ratios. The range of concentration used were 0.05 – 0.54N for individual KOH and NaOH washing solutions and between 0.054N to 0.216N for the mix solution. Their effects on frying of treated breaded fillets on the rancidity development and physico-chemical changes in refined bleached and deodorized (RBD) palm olein, the frying oil used was also carried out. Washing treatment with either KOH or NaOH solution was found to affect the rancidity development of the muscle. PV and TBARS values were significantly increased ($p < 0.05$) in all treatments. The absorbance for hydroperoxides at 1745 cm^{-1} using Fourier Transform Infrared (FTIR) spectroscopy for oil extracted from all treated fillets also increased. The correlation coefficients (R^2) between absorbance versus PV for KOH and NaOH were 0.8467 and 0.6651, respectively. Fatty acids composition and

fatty acids ratios ($C_{18:1}+C_{18:2}$)/ $C_{16:0}$ and $C_{18:2}/C_{16:0}$ showed a decreasing trend in total unsaturated fatty acids with increasing concentration of washing solutions. Nineteen triglycerides were determined and only four were detected in the samples. The percentage of tripetroselinin and triolein decreased while trilaurin increased with increasing concentration of both KOH and NaOH. Sensory evaluation for odor, texture and acceptability indicated that panelists have more preferences for samples treated with less than 0.36N for both alkalies. The rancid odor was detected at TBARS value of 17 μmol malonaldehyde/kg and the PV at 20 meq hydroperoxide/kg oil. PV was found to be highest in samples washed in mixture of 0.108N KOH and 0.162N NaOH. TBARS and FFA values were highest in samples washed in mixture of 0.162N KOH and 0.108N NaOH and mixture of 0.108N KOH and 0.162N NaOH, respectively. Nonetheless, sensory evaluation indicated no significant difference ($p>0.05$) among all samples. There were significant differences ($p<0.05$) in PV, IV, AnV and totox values between control and treatments during 5 consecutive days of frying. However, there was no significant difference ($p>0.05$) in FFA and ratio of $C_{18:2}/C_{16:0}$ between control and treatments during frying. Instrumental color determination showed increases in red (R) and yellow (Y) values of the frying oil, thus, indicating darkening of the oil. For fried breaded fillets, TBARS values increased and significant differences ($p<0.05$) were obtained between control and treatments. No significant different ($p>0.05$) in organoleptic quality of fried fillets between control and treatments. The amount of alkaline contaminant materials (ACM) were significantly ($p<0.05$) increased during 5 consecutive days of frying. Washing treatments did not seem to affect the ACM content.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**KESAN PEMBASUHAN MENGGUNAKAN CAMPURAN KALIUM DAN
NATRIUM HIDROKSIDA KE ATAS PERUBAHAN KETENGIKAN DALAM
FILET IKAN TILAPIA (*OREOCHROMIS MOSSAMBICUS*) DAN MINYAK
PENGGORENGAN**

Oleh

ANIDA YUSOFF

Januari 2003

Pengerusi : Professor Madya Dr. Jamilah Bakar, Ph.D.

Fakulti : Sains Makanan dan Bioteknologi

Satu kajian telah dijalankan untuk menyelidik keberkesanan penggunaan kalium (KOH) dan natrium hidroksida (NaOH) secara individu dan campuran keduanya dalam penyahbauan dan citarasa lumpur di dalam filet ikan tilapia (*Oreochromis mossambicus*). Empat larutan campuran KOH dan NaOH digunakan pada nisbah yang berlainan. Kepekatan yang digunakan adalah antara 0.05 – 0.54N bagi setiap larutan individu dan bagi larutan campuran, kepekatan adalah antara 0.054N dan 0.216N. Kesan rawatan terhadap perubahan ketengikan dan fiziko kimia di dalam filet goreng bersalut dan minyak gorengan yang digunakan juga telah dijalankan. Kaedah pembasuhan sama ada menggunakan KOH atau NaOH telah menunjukkan kesan terhadap ketengikan dalam isi ikan. Peningkatan yang ketara ($p < 0.05$) bagi nilai peroksida (PV) dan asid tiobarbiturik (TBARS) telah diperolehi dalam semua rawatan. Nilai penyerapan hidroperoksida pada 1745 cm^{-1} menggunakan Fourier Transform Infrared (FTIR) spektroskopi bagi minyak yang diekstrak dari filet yang dirawat juga

meningkat. Pekali perkaitan (R^2) bagi nilai penyerapan melawan PV adalah 0.8467 bagi larutan KOH dan 0.6651 bagi larutan NaOH. Komposisi asid lemak dan nisbah asid lemak ($C_{18:1}+C_{18:2}/C_{16:0}$) dan ($C_{18:2}/C_{16:0}$) telah menunjukkan nilai yang menurun bagi jumlah asid lemak tak tepu dengan peningkatan kepekatan larutan pembasuh. Sembilan belas trigliserida telah diuji tetapi hanya empat dapat dikesan di dalam sampel. Peratus tripetroselinin dan triolein berkurang dengan peningkatan kepekatan larutan pembasuh. Manakala, trilaurin telah bertambah bagi kedua-dua larutan pembasuh. Ujian penilaian deria bagi bau, tekstur dan penerimaan di atas filet telah menunjukkan bahawa ahli panel menyukai sampel yang dirawat dengan alkali pada kepekatan kurang dari 0.36N bagi kedua-dua NaOH dan KOH. Bau ketengikan diperolehi pada nilai TBARS 17 μmol MA/kg dan PV pada 20 meq hidroperoksida/kg minyak. PV adalah tertinggi bagi campuran basuhan 0.108N KOH dan 0.162N NaOH. Nilai TBARS dan asid lemak bebas (FFA) adalah tertinggi dalam sampel yang dibasuh dengan larutan campuran 0.162N KOH dan 0.108N NaOH dan larutan campuran 0.108N KOH dan 0.162N NaOH. Bagaimanapun, tiada perbezaan yang ketara ($p>0.05$) diperolehi dalam semua sampel bagi penilaian deria. Terdapat perbezaan yang ketara ($p<0.05$) dalam PV, nilai iodin (IV), nilai anisidin (AnV) dan nilai totox antara kawalan dan sampel rawatan sepanjang 5 hari penggorengan berturut-turut. Tiada perbezaan yang ketara ($p>0.05$) dalam FFA dan nisbah asid lemak ($C_{18:2}/C_{16:0}$) antara kawalan dan sampel rawatan semasa penggorengan. Pengukuran warna menunjukkan peningkatan dalam nilai merah (R) dan kuning (Y) bagi minyak goreng yang menandakan berlaku penghitaman dalam warna minyak kelapa sawit semasa penggorengan. Bagi filet bersalut yang digoreng, nilai TBARS menunjukkan peningkatan dan perbezaan yang ketara ($p<0.05$) di antara

kawalan dan sampel rawatan. Tiada perbezaan yang ketara ($p>0.05$) antara kawalan dan sampel rawatan dari aspek deria bagi gorengan filet bersalut. Kandungan kontaminasi alkali (ACM) dalam minyak menunjukkan peningkatan yang ketara ($p<0.05$) sepanjang 5 hari penggorengan berturut-turut bagi semua minyak goreng, rawatan pembasuhan tidak memberi kesan ke atas kandungan ACM.

ACKNOWLEDGEMENTS

Alhamdulillah, first of all I would like to express my utmost thanks and gratitude to Almighty Allah S.W.T. who has given me the will, strength and patience to complete this project and my *Salawat* and *Salam* to His righteous messenger, Prophet Muhammad S.A.W.

I would like to express my sincere appreciation to Associate Professor Dr. Hjh. Jamilah Bakar, the Chairman of my Supervisory Committee, of the Department of Food Technology, Faculty of Food Science and Biotechnology for her guidance, patience, understanding, encouragement and supervisions throughout the course of study till the preparation of this thesis. I am also very grateful to other members of the supervisory committee, Prof. Dr. Yaakob B. Che Man and Prof. Dr. Jinap Selamat of the Faculty Food Science and Biotechnology, Universiti Putra Malaysia for their supervision, support and comments.

I would like to acknowledge the financial support provided by the IRPA fund for the research performed at UPM, and by the Universiti Teknologi Mara (UiTM) for awarding me the Tenaga Pengajar Muda (TPM) scholarship during my study at UPM.

My sincere appreciation is also to all the technical staffs, graduate students and research assistants in the Faculty of Food Science and Biotechnology for their co-operation, moral encouragement and support to complete this study.

Not the least but the most important, my deepest appreciation to my beloved parents, sisters, twin sister and brother for their moral support and understanding throughout my studies. I shall forever recognize for their sacrifices they have given in anyway during the many years of my pursue for knowledge. I wish will be in Allah's baraqah.



I certify that an Examination Committee met on 14th January 2003 to conduct the final examination of Anida Yusoff on her Master of Science thesis entitled “Effects of Washing with Potassium and Sodium Hydroxide Mixtures in the Development of Rancidity in Black Tilapia (*Oreochromis mossambicus*) Fillets and the Frying Oil” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulation 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows :

Salmah Yusof, Ph.D.

Associate Professor,
Department of Food Technology,
Faculty of Food Science and Biotechnology,
Universiti Putra Malaysia.
(Chairperson)

Jamilah Bakar, Ph.D.

Associate Professor,
Department of Food Technology,
Faculty of Food Science and Biotechnology,
Universiti Putra Malaysia.
(Member)

Yaakob B. Che Man, Ph.D.

Professor,
Department of Food Technology,
Faculty of Food Science and Biotechnology,
Universiti Putra Malaysia.
(Member)

Jinap Selamat, Ph.D.

Professor,
Department of Food Science,
Faculty of Food Science and Biotechnology,
Universiti Putra Malaysia.
(Member)



GULAM RUSUL RAHMAT ALI, Ph.D.

Professor / Deputy Dean,
School of Graduate Studies,
Universiti Putra Malaysia.

Date : 25 MAR 2003

This thesis submitted to the Senate of Universiti Putra Malaysia and was accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

Jamilah Bakar, Ph.D.

Associate Professor,
Department of Food Technology,
Faculty of Food Science and Biotechnology,
Universiti Putra Malaysia.
(Chairperson)

Yaakob B. Che Man, Ph.D.

Professor,
Department of Food Technology,
Faculty of Food Science and Biotechnology,
Universiti Putra Malaysia.
(Member)

Jinap Selamat, Ph.D.

Professor,
Department of Food Science,
Faculty of Food Science and Biotechnology,
Universiti Putra Malaysia.
(Member)



AINI IDERIS, Ph.D.

Professor / Dean,
School of Graduate Studies
Universiti Putra Malaysia

Date: 10 APR 2003

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledge. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



ANIDA BINTI YUSOFF

Date : 3rd March 2003

TABLE OF CONTENTS

	Page
DEDICATION.....	ii
ABSTRACT.....	iii
ABSTRAK.....	v
ACKNOWLEDGEMENTS.....	viii
APPROVAL SHEET.....	x
DECLARATION FORM.....	xii
LIST OF TABLES.....	xvii
LIST OF FIGURES.....	xix
LIST OF PLATES.....	xxi
LIST OF ABBREVIATIONS.....	xxii
 CHAPTER	
I GENERAL INTRODUCTION.....	1
II LITERATURE REVIEW.....	4
Commercial Production of Tilapia.....	4
Factors Affecting the Acceptance of Tilapia.....	4
Musty-Muddy Off-odor and Flavors in Freshwater Fish.....	6
Techniques of Removing Muddy Off-odor and Flavor....	8
Effects of Organic Acid and Alkaline Washing on the Quality of Fish.....	11
Physical and Chemical Changes.....	11
Microbiological Changes.....	13
Sensory Changes.....	13
Mechanism of Alkaline Washing on Oxidation Deterioration.....	14
Contributing Factors in the Effectiveness of Alkali in Removing Muddy Off-odor and Flavor.....	15
Solubility of NaOH and KOH in Water.....	15
Ionic Strength.....	16
Ph of Medium.....	17
The Lipid Rancidity in Freshwater and Marine Fish.....	18
Distribution of Fatty acids Composition in Fish Muscle.....	21
Classes of Lipids in Fish.....	21
Extraction of lipid from Fish Muscle.....	22
Lipid Oxidation and Its Implication.....	24



General Deterioration Reactions.....	25
Oxidative Rancidity in Fish Muscle.....	27
Effects of Frying on Quality of Oil and Breaded Fillets...	28
Reactions during Frying.....	30
Measurement of Rancidity in Fish Lipid and Frying Oils.	32
Chemical Based Analyses.....	32
Instrumental Analyses.....	35
Fourier Transform Infrared (FTIR) Spectroscopy	
Analysis.....	35
Sensory Evaluation.....	36
 III EFFECTS OF INDIVIDUAL POTASSIUM AND	
SODIUM HYDROXIDE WASHING ON	
RANCIDITY DEVELOPMENT IN BLACK	
TILAPIA (<i>Oreochromis mossambicus</i>)	
MUSCLE.....	39
Introduction.....	39
Materials and Methods.....	42
Washing Treatment.....	42
Lipid Extraction.....	43
Chemical Analysis.....	45
pH Measurement.....	45
Peroxide Values (PV).....	45
Thiobarbituric Acid Reactive Substances (TBARS)	
Values.....	46
Fatty Acids Composition.....	48
Ratio of fatty acids $C_{18:2}/C_{16:0}$ and $C_{18:1}+C_{18:2}/C_{16:0}$	48
Triglycerides (TAG) Analysis by HPLC.....	49
Fourier Transform Infrared (FTIR) Spectroscopy of	
Extracted Lipid.....	50
Sensory Evaluation.....	51
Statistical Analysis.....	52
Results and Discussion.....	52
Chemical Analysis.....	52
pH Values.....	52
Peroxide Values (PV).....	54
TBARS Values.....	55
Fatty Acids Composition.....	59
Ratio of Fatty Acids($C_{18:2}/C_{16:0}$).....	62
Triglycerides Profile.....	67
FTIR-based Analysis.....	73
Sensory Evaluation.....	80
Conclusion.....	86
 IV EFFECTS OF POTASSIUM AND SODIUM	

HYDROXIDE WASHING MIXTURE ON RANCIDITY DEVELOPMENT IN BLACK TILAPIA (*Oreochromis mossambicus*) MUSCLE..... 88

Introduction.....	88
Materials and Methods.....	90
Washing Treatment.....	90
Lipid Extraction.....	90
Chemical Analysis.....	91
Free Fatty Acid(FFA).....	91
Sensory Evaluation.....	92
Statistical Analysis.....	92
Results and Discussion.....	93
pH	93
PV.....	94
TBARS	95
FFA.....	95
Fatty Acids Ratio ($C_{18:2}/C_{16:0}$).....	96
Sensory Evaluation.....	98
Conclusion.....	101

V PHYSICOCHEMICAL CHANGES IN RBD PALM OIL DURING FRYING AND THE QUALITY OF FRIED BREADED TREATED FILLETS..... 103

Introduction.....	103
Materials and Methods.....	105
Fish Fillet Preparation.....	105
Washing Treatment.....	105
Battering and Breading.....	106
Frying Experiments.....	106
Chemical Analysis of Fried RBD Palm Olein.....	107
Physical Analysis of Fried RBD Palm Olein.....	109
Color Determination by Lovibond Tintometer.....	109
Chemical Analysis of Fried Fillets.....	109
TBARS Values.....	109
Alkaline Contaminant Materials (ACM) Quick Test Method.....	109
Preparation of Test Solution.....	109
Preparation of Standard Curve.....	110
Determination of Standard Solution by Hunterlab Colorimeter.....	111
Sensory Evaluation of Fried Battered Fillets.....	111
Statistical Analysis.....	112
Results and Discussion.....	113
Chemical Analysis of Fried RBD Palm Olein.....	113

Changes in PV, AnV and IV During Frying.....	113
Changes in Totox Values, FFA and ($C_{18:2}/C_{16:0}$) Ratio.....	117
Fatty Acids Composition.....	120
ACM Quick Test Method.....	122
Physical Analysis of Fried RBD Palm Olein.....	125
Changes in the Color of RBD Palm Olein During Frying.....	125
Chemical Analysis of Fried Fillets.....	129
TBARS Values of Fried Breaded Fillets.....	129
Sensory Evaluation of Fried Breaded Fillets.....	131
Conclusion.....	141
VI SUMMARY AND CONCLUSION.....	142
REFERENCES.....	145
APPENDICES.....	164
BIODATA OF THE AUTHOR.....	178

LIST OF TABLES

Table		Page
1	The solubility of NaOH and KOH in Water at a temperature range of 0 to 30°C.....	17
2	Principal pathways for formation of frying fat alteration compounds.....	31
3	Fatty acids composition of extracted fish oil from treated fillets with increasing concentration of washing solution KOH and NaOH.....	61
4	Total saturated, unsaturated fatty acid and ratio $C_{18:1}+C_{18:2}/C_{16:0}$ during washing with KOH and NaOH solution.....	65
5	Changes in the ratio of fatty acids $C_{18:2}/C_{16:0}$ during washing with KOH and NaOH.....	66
6	Retention time (RT) of Triglycerides Standard.....	70
7	Triglycerides (TAG) composition (%) of extracted fish oil from treated fillets with increasing concentration of washing solution KOH.....	71
8	Triglycerides (TAG) composition (%) of extracted fish oil from treated fillets with increasing concentration of washing solution NaOH.....	72
9	Sensory scores of uncooked black tilapia fillet washed at different concentration of KOH and NaOH washing solution	83
10	The washing solutions consisting of KOH and NaOH mixture at different concentrations and ratios.....	91
11	The effect of washing treatments on the pH, PV, TBARS values, ratios $C_{18:2}/C_{16:0}$ and Free fatty acid of extracted fish oil from treated fillets with different mixtures of potassium and sodium hydroxide washing solutions.....	97
12	Sensory scores of uncooked black tilapia fillet washed with mixtures of KOH and NaOH washing	



	solutions.....	100
13	Changes in PV, AnV and IV of RBD palm olein during frying of breaded fillet washed with 0.05N KOH and 0.05N NaOH.....	116
14	Changes in Totox values, free fatty acids (FFA) content and C _{18:2} /C _{16:0} ratio of RBD palm olein during frying of breaded fillet washed with 0.05N KOH and 0.05N NaOH	119
15	Fatty acids composition for control, 0.05N KOH and 0.05N NaOH of RBD palm olein during frying.....	121
16	Concentration of alkaline contaminant material (ACM), mg/100g in RBD palm olein during intermittent frying of treated fillets.....	124
17	Changes in color of RBD palm olein during frying of breaded fillets washed with 0.05N KOH and 0.05N NaOH.....	126
18	Changes in TBARS values of breaded fillets fried in RBD palm olein.....	130
19	Mean sensory scores of fried breaded fillets of treated black tilapia (<i>Oreochromis mossambicus</i>) in RBD palm olein for different days of frying.....	134
20	Changes in pH during washing with KOH and NaOH.....	165
21	Changes in peroxide values during washing with KOH and NaOH.....	166
22	Changes in TBARS values during washing with KOH and NaOH.....	167
23	Changes of absorbance at 1745 cm ⁻¹ of extracted oil and PV for KOH and NaOH treated samples.....	168
24	Color changes of alkaline contaminant material (ACM) in test solution.....	173
25	The effect of washing in different mixtures of potassium and sodium hydroxide washing solutions on the pH, PV, TBARS values, Fatty acids ratio (C _{18:2} /C _{16:0}) and Free Fatty Acids of extracted oil from samples.....	177

LIST OF FIGURES

Figure		Page
1	The free-radical chain mechanism.....	24
2	Some routes of decomposition of fat hydroperoxide.....	26
3	Changes in pH values of fillets after washing with different concentrations of KOH and NaOH.....	56
4	Changes in Peroxide values (PV) of extracted oil from treated fillet with increasing concentration of washing solution KOH and NaOH.....	57
5	Changes in TBARS values of extracted oil from treated fillet with increasing concentration of washing potassium hydroxide (KOH) and sodium hydroxide (NaOH).....	58
6	Fourier Transform Infrared (FTIR) spectra of black tilapia extracted oil obtained in the 4000-400 cm^{-1} for various concentration of KOH washing treatments	76
7	Fourier Transform Infrared (FTIR) spectra of black tilapia extracted oil obtained in the 4000-400 cm^{-1} for various concentration of NaOH washing treatments	77
8	Differential spectra of black tilapia extracted oil obtained at 1745.00 cm^{-1} for various concentration of KOH washing treatments.....	78
9	Plot of absorbance at 1745.00 cm^{-1} of extracted oil versus PV for KOH treated samples.....	78
10	Differential spectra of black tilapia extracted oil obtained at 1745.00 cm^{-1} for various concentration of NaOH washing treatments.....	79
11	Plot of absorbance at 1745.00 cm^{-1} of extracted oil versus PV for NaOH treated samples.....	79
12	Flow-chart for frying process of treated fish fillets.....	108
13	Standard curve for different concentration of TEP.....	164



14	Spectrum from various concentration of sodium palmitate measured at wavelengths 460 nm to 560 nm.....	174
15	A linear regression plot of absorbances at 540 nm vs concentration of sodium palmitate in fresh RBD palm olein.....	174
16	HPLC Chromatogram of 0.05N KOH (A) and 0.09N KOH (B) Washing Solution (Abbreviations : Peak 1: Triolein; Peak 2: Tripetroselinin; Peak 3 : Unknown; Peak 4 : Unknown; Peak 5 : Trilaurin).....	175
17	HPLC Chromatogram of Control (A) and 0.05N NaOH (B) Washing Solution (Abbreviations : Peak 1: Triolein; Peak 2: Tripetroselinin; Peak 3 : Unknown; Peak 4 : Unknown; Peak 5 : Trilaurin).....	176

LIST OF PLATES

Plate		Page
1	Wild black tilapia (<i>Oreochromis mossambicus</i>) and the fillet.....	44
2	Appearance of treated black tilapia fillets washed with 0.0, 0.5, 1.0, 1.5, 2.0, 2.5 and 3.0% which is equivalent to 0.00N, 0.09N, 0.18N, 0.27N, 0.36N, 0.45N and 0.54N of potassium hydroxide (KOH) washing solution.....	84
3	Appearance of treated black tilapia fillets washed with 0.0, 0.5, 1.0, 1.5, 2.0, 2.5 and 3.0% which is equivalent to 0.00N, 0.09N, 0.18N, 0.27N, 0.36N, 0.45N and 0.54N of sodium hydroxide (NaOH) washing solution.....	85
4	Changes in color of RBD palm olein during frying of 5 consecutive days of control breaded fillet.....	127
5	Changes in color of RBD palm olein during frying of 5 consecutive days for breaded fillet washed with 0.05N KOH washing solution.....	127
6	Changes in color of RBD palm olein during frying of 5 consecutive days for breaded fillet washed with 0.05N NaOH washing solution.....	128
7	Overall changes in color of RBD palm olein during of 5 consecutive days for control, 0.05N KOH and 0.05N NaOH washed fillet respectively.....	128
8	Fried black tilapia breaded fillet of control at day 5 of frying.....	139
9	Fried black tilapia breaded fillet of 0.05N KOH at day 5 of frying.....	139
10	Fried black tilapia breaded fillet of 0.05N NaOH at day 5 of frying.....	140



LIST OF ABBREVIATIONS

ACM	-	Alkaline contaminant material
ACS	-	American Chemical Society
ANOVA	-	Analysis of Variance
AnV	-	Anisidine Value
cm	-	centimeter
FFA	-	Free Fatty Acid
FTIR	-	Fourier Transform Infra-red
g	-	gram
i.d	-	inner diameter
IV	-	Iodine Value
KOH	-	Potassium hydroxide
Meq	-	milliequivalent
mL	-	milliliter
N	-	normality
NaOH	-	Sodium hydroxide
nm	-	nanometer
PV	-	Peroxide Value
RBD	-	Refined, Bleached and Deodorized
rpm	-	revolution per minute
SAS	-	Statistical Analysis System
TBARS	-	Thiobarbituric Acid Reactive Substances
w/v	-	weight/volume

% - percentage

μ - micro

CHAPTER I

GENERAL INTRODUCTION

Depletion of many ocean fish species has resulted in the use of lesser known fish (NAS, 1978). However, successful marketing of those fish for commercial purposes depends on consumer acceptance of their sensory properties such as flavor and textural characteristics. Flavor and flavor characteristics are concluded as the most important determinant of consumer acceptability (Connell and Howgate, 1971; Hamilton and Bennett, 1983 & 1984; Wesson *et al.*, 1979; Sawyer *et al.*, 1988). However, for fish with mild flavor, i.e those which exhibited low to moderate intensity of fishy and/or oxidized flavors, texture is more important (Wesson *et al.*, 1979; Sawyer *et al.*, 1988). A serious and frequent problem in intensive fish culture is the absorption of objectionable flavor compounds by fish from the culture environment which will contribute to the earthy odor and flavor (Johnsen, 1989). The fish readily absorb organic compounds through their gills and skin as well as from the digestive tract.

Many studies have been conducted to remove the earthy odor and flavor in fish. The treatments for removing muddy flavor and odor of black tilapia involved chemical and physiological approaches. Several traditional methods which assist in removing the muddy flavor and odor in tilapia include the washing with tamarind pulp and lemon juice or mixture of tamarind pulp, *Garcinia pyrifera* (asam gelugor), guava leaves, flour, “lambanog”, salt and lemon grass (Jamilah and Siti Aini, 1997; Mabesa *et al.*, 1989; Rohani and Yunus, 1994; Anon, 1991). However,